

# *Diplotaxis virgata*

## Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **8.1**

Temperature indicator value: **8.3**

Moisture indicator value: **3**

Reaction indicator value: **6.7**

Nutrient indicator value: **5.1**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.75**

Disturbance frequency (herb layer): **2.07**

Disturbance severity: **0.78**

Disturbance severity (herb layer): **0.77**

Mowing frequency: **0.54**

Grazing pressure: **0.24**

Soil disturbance: **0.45**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**