# *Doronicum columnae*

## Habitus and growth type

Plant height [m]: **0.32**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.54**
Dispersal mode: **Anemochory**
Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **3**
Moisture indicator value: **6**
Reaction indicator value: **7**
Nutrient indicator value: **6.5**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.17**
Disturbance frequency (herb layer): **1.1**
Disturbance severity: **0.57**
Disturbance severity (herb layer): **0.16**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0.02**
Grazing pressure: **0.19**
Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae*, CS (MUL) *Mulgedio-Aconitetea*, KE (THL) *Thlaspietea rotundifolii***

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **5**
Continentality amplitude: **1**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*