

# ***Doronicum plantagineum***

## **Habitus and growth type**

Plant height [m]: **0.55**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## **Flower**

Flowering period: **April-July**

## **Fruit, seed and dispersal**

Dispersal mode: **Anemochory, Anthropochory**

Dispersal distance class: **3, 7**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **6.3**

Moisture indicator value: **6**

Reaction indicator value: **6.7**

Nutrient indicator value: **4.7**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.05**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.67**

Disturbance severity (herb layer): **0.12**

Mowing frequency: **0.02**

Grazing pressure: **0.22**

Soil disturbance: **0.09**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) Quercetea ilicis](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**