

Dysphania botrys

Habitus and growth type

Plant height [m]: **0.4**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.19**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Reaction indicator value: **6.6**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.61**

Disturbance frequency (herb layer): **1.7**

Disturbance severity: **0.77**

Disturbance severity (herb layer): **0.77**

Mowing frequency: **0.31**

Grazing pressure: **0.15**

Soil disturbance: **0.52**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QD \(DIG\) Digitario sanguinalis-Eragrostietea minoris](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region](#)

Constant species of EUNIS habitats: [U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region](#)

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: 7

Continentality amplitude: 9