

# *Elatine triandra*

## Habitus and growth type

Plant height [m]: **0.04**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **May-September**

## Fruit, seed and dispersal

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **6.3**

Moisture indicator value: **9**

Reaction indicator value: **4.4**

Nutrient indicator value: **4.6**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.91**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.71**

Disturbance severity (herb layer): **0.71**

Mowing frequency: **0.01**

Grazing pressure: **0.16**

Soil disturbance: **0.37**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q62 Periodically exposed shore with stable, mesotrophic sediments with pioneer or ephemeral vegetation](#), [V14 Inundated or inundatable cropland, including rice fields](#)

Constant species of EUNIS habitats: [V14 Inundated or inundatable cropland, including rice fields](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Synanthropic**

## Distribution

Continentality: **5**

Continentality amplitude: **4**