

Eleocharis acicularis

Habitus and growth type

Plant height [m]: **0.08**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte, Hydrophyte**

Leaf

Specific leaf area [mm²/mg]: **77.33**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.1**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**

Temperature indicator value: **5.6**

Moisture indicator value: **10.1**

Nutrient indicator value: **2.6**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.67**

Disturbance frequency (herb layer): **1.7**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.08**

Grazing pressure: **0.13**

Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **[OC \(ISO\) Isoëto-Nanojuncetea](#)**

EUNIS habitat

Diagnostic species of EUNIS habitats: **[Q62 Periodically exposed shore with stable, mesotrophic sediments with pioneer or ephemeral vegetation](#), [V14 Inundated or inundatable cropland, including rice fields](#)**

Constant species of EUNIS habitats: **[Q62 Periodically exposed shore with stable, mesotrophic sediments with pioneer or ephemeral vegetation](#)**

Dominant species of EUNIS habitats: **[Q62 Periodically exposed shore with stable, mesotrophic sediments with pioneer or ephemeral vegetation](#)**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **9**