# *Epilobium fleischeri*

### Habitus and growth type

Plant height [m]: **0.26**
Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

### Flower

Flowering period: **May-August**

### Fruit, seed and dispersal

Seed mass [mg]: **0.18**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **3.5**
Moisture indicator value: **4**
Reaction indicator value: **2**
Nutrient indicator value: **2**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **1.1**
Disturbance frequency (herb layer): **1.44**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.46**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0.02**
Grazing pressure: **0.17**
Soil disturbance: **0.32**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **KE (THL) *Thlaspietea rotundifolii***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **U71 Unvegetated or sparsely vegetated gravel bar in montane and alpine regions**
Constant species of EUNIS habitats: **U71 Unvegetated or sparsely vegetated gravel bar in montane and alpine regions**

### Broad habitat

Occurrence in broad habitats: **Spring, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **5**
Continentality amplitude: **3**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*