

Equisetum scirpoides

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Moisture indicator value: **6**

Disturbance indicator values

Disturbance frequency: **0.78**

Disturbance frequency (herb layer): **1.45**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.01**

Grazing pressure: **0.24**

Soil disturbance: **0.09**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [U35 Boreal and Arctic base-rich inland cliff](#)

Constant species of EUNIS habitats: [T3H Larix light taiga](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Forest, Sparsely vegetated (incl. rock and scree)**