Equisetum telmateia

Habitus and growth type

Plant height [m]: **1.22**
Life span: **Perennial**
Life form: **Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **16.26**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.9**
Moisture indicator value: **7.8**
Reaction indicator value: **7.6**
Nutrient indicator value: **5.4**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.45**
Disturbance frequency (herb layer): **1.44**
Disturbance severity: **0.59**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.38**
Mowing frequency: **0.24**
Grazing pressure: **0.18**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **HA (POP) *Alno glutinosae-Populetea albae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T14 Mediterranean and Macaronesian riparian forest**
Constant species of EUNIS habitats: **T14 Mediterranean and Macaronesian riparian forest**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Forest, Synanthropic**

Distribution

Continentality: **3**
Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz