

Equisetum variegatum

Habitus and growth type

Plant height [m]: **0.28**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **6.33**

Flower

Flowering period: **April-September**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **3.6**

Moisture indicator value: **8.4**

Reaction indicator value: **7.4**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.28**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.28**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.27**

Grazing pressure: **0.15**

Soil disturbance: **0.07**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [PA \(SCH\) *Scheuchzerio palustris-Caricetea fuscae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q45 Arctic-alpine rich fen](#), [Q46 Carpathian travertine fen with halophytes](#)

Constant species of EUNIS habitats: [Q41 Alkaline, calcareous, carbonate-rich small-sedge spring fen](#), [Q45 Arctic-alpine rich fen](#), [Q46 Carpathian travertine fen with halophytes](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Mire, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **5**