Eremopyrum triticeum

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **5.3**
Moisture indicator value: **3**

Disturbance indicator values

Disturbance frequency: **2.01**
Disturbance frequency (herb layer): **2.1**
Disturbance severity: **0.49**
Disturbance severity (herb layer): **0.48**
Mowing frequency: **0.13**
Grazing pressure: **0.37**
Soil disturbance: **0.37**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DB (FEP) *Festuco-Puccinellietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R64 Semi-desert salt pan**, **S67 Aralo-Caspian semi-desert**, **S68 Semi-desert sand dune with sparse scrub**, **S94 Semi-desert riparian scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Constant species of EUNIS habitats: **R1C Desert steppe**, **R64 Semi-desert salt pan**, **R65 Continental subsaline alluvial pasture and meadow**, **S67 Aralo-Caspian semi-desert**, **S68 Semi-desert sand dune with sparse scrub**, **S94 Semi-desert riparian scrub**

Dominant species of EUNIS habitats: **S68 Semi-desert sand dune with sparse scrub**

Broad habitat

Occurrence in broad habitats: **Saline vegetation, Semi-desert, Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz