Erica multiflora

Habitus and growth type

Plant height [m]: **0.98**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **4.39**

Flower

Flowering period: **August-November**

Fruit, seed and dispersal

Seed mass [mg]: **0.08**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **8.7**
Moisture indicator value: **2.5**
Reaction indicator value: **6.5**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.34**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.23**
Mowing frequency: **0.02**
Grazing pressure: **0.31**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S54 Thermomediterranean arid scrub**, **S61 Western basiphilous garrigue**
Constant species of EUNIS habitats: **S54 Thermomediterranean arid scrub**, **S61 Western basiphilous garrigue**
Dominant species of EUNIS habitats: **S61 Western basiphilous garrigue**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz