Erodium foetidum

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.6**
Temperature indicator value: **5.1**
Moisture indicator value: **3.2**
Reaction indicator value: **6.2**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.11**
Disturbance frequency (herb layer): **1.15**
Disturbance severity: **0.38**
Disturbance severity (herb layer): **0.33**
Mowing frequency: **0.13**
Grazing pressure: **0.31**
Soil disturbance: **0.12**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1H Iberian oromediterranean basiphilous dry grassland**, **S73 Western Mediterranean mountain hedgehog-heath**, **U34 Mediterranean siliceous inland cliff**

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz