

# *Euphorbia amygdaloides*

## Habitus and growth type

Plant height [m]: **0.44**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **24.04**

## Flower

Flowering period: **January-June**

## Fruit, seed and dispersal

Seed mass [mg]: **4.17**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Moisture indicator value: **5.3**

Reaction indicator value: **6.7**

Nutrient indicator value: **5.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.04**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.13**

Mowing frequency: **0.01**

Grazing pressure: **0.2**

Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) \*Carpino-Fagetea sylvaticae\*](#), [CB \(PUB\) \*Quercetea pubescentis\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T17 \*Fagus forest on non-acid soils\*](#), [T1G \*Alnus cordata forest\*](#), [T39 \*Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest\*](#), [T3C \*Taxus baccata forest\*](#)

Constant species of EUNIS habitats: [S37 \*Corylus avellana scrub\*](#), [T17 \*Fagus forest on non-acid soils\*](#), [T19 \*Temperate and submediterranean thermophilous deciduous forest\*](#), [T1E \*Carpinus and Quercus mesic deciduous forest\*](#), [T1G \*Alnus cordata forest\*](#), [T32 \*Temperate mountain Abies forest\*](#), [T33 \*Mediterranean mountain Abies forest\*](#), [T39 \*Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest\*](#), [T3C \*Taxus baccata forest\*](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**

Continentality amplitude: **4**