

Euphorbia glabriflora

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **6.7**

Moisture indicator value: **3.7**

Reaction indicator value: **7.1**

Nutrient indicator value: **2.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.99**

Disturbance frequency (herb layer): **1.61**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.23**

Grazing pressure: **0.29**

Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) *Sedo-Scleranthetea*](#), [DA \(FES\) *Festuco-Brometea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the](#)

[Balkans, U3A Temperate ultramafic inland cliff](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Dominant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**