

# *Festuca beckeri*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.64**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **4.5**

Moisture indicator value: **4.8**

Disturbance indicator values

Disturbance frequency: **1.29**

Disturbance frequency (herb layer): **1.65**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.24**

Grazing pressure: **0.17**

Soil disturbance: **0.39**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [BD \(PYR\) \*Pyrolo-Pinetea sylvestris\*](#),  
[CJ \(COR\) \*Koelerio-Corynephoretea canescentis\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [N17 Black Sea coastal dune grassland \(grey\)](#)

**dune), R11 Pannonian and Pontic sandy steppe**

Constant species of EUNIS habitats: **N17 Black Sea coastal dune grassland (grey dune), R11 Pannonian and Pontic sandy steppe**

Dominant species of EUNIS habitats: **R11 Pannonian and Pontic sandy steppe**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Forest**