

# *Festuca indigesta*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **3**

Moisture indicator value: **2**

Reaction indicator value: **3**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.1**

Disturbance frequency (herb layer): **1.73**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.1**

Grazing pressure: **0.31**

Soil disturbance: **0.12**

## Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1G Iberian oromediterranean siliceous dry grassland](#), [R1N Open Iberian supramediterranean dry acid and neutral grassland](#)

Constant species of EUNIS habitats: [R1G Iberian oromediterranean siliceous dry grassland](#), [R1N Open Iberian supramediterranean dry acid and neutral grassland](#)

Dominant species of EUNIS habitats: [R1G Iberian oromediterranean siliceous dry](#)

**[grassland, R1N Open Iberian supramediterranean dry acid and neutral grassland](#)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub**