

# *Filago arvensis*

## Habitus and growth type

Plant height [m]: **0.2**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **39.06**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.21**

Dispersal mode: **Anemochory, Anthropochory**

Dispersal distance class: **5, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **7.3**

Moisture indicator value: **2.6**

Reaction indicator value: **4.5**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.32**

Disturbance frequency (herb layer): **1.96**  
Disturbance severity: **0.57**  
Disturbance severity (herb layer): **0.55**  
Mowing frequency: **0.54**  
Grazing pressure: **0.31**  
Soil disturbance: **0.37**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) \*Koelerio-Coryneporetea canescentis\*](#), [QB \(SIS\) \*Sisymbrietea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#)

Constant species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

## Distribution

Continentality: **7**  
Continentality amplitude: **8**