

Filipendula ulmaria

Habitus and growth type

Plant height [m]: **0.92**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **28.99**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.68**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**

Temperature indicator value: **4.9**

Moisture indicator value: **7.9**

Reaction indicator value: **5.8**

Nutrient indicator value: **5**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.66**

Disturbance frequency (herb layer): **1.54**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.44**

Grazing pressure: **0.18**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CM \(MOL\) Molinio-Arrhenatheretea**](#),
[**IA \(ALN\) Alnetea glutinosae**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R35 Moist or wet mesotrophic to eutrophic hay meadow**](#), [**T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**](#)

Constant species of EUNIS habitats: [**Q42 Extremely rich moss-sedge fen**](#), [**Q53 Tall-sedge bed**](#), [**R22 Low and medium altitude hay meadow**](#), [**R35 Moist or wet mesotrophic to eutrophic hay meadow**](#), [**R37 Temperate and boreal moist or wet oligotrophic grassland**](#), [**R55 Lowland moist or wet tall-herb and fern fringe**](#), [**S37 Corylus avellana scrub**](#), [**S92 Salix fen scrub**](#), [**T11 Temperate Salix and Populus riparian forest**](#), [**T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**](#), [**T13 Temperate hardwood riparian forest**](#), [**T15 Broadleaved swamp forest on non-acid peat**](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality amplitude: **6**