

# *Fourraea alpina*

## Habitus and growth type

Plant height [m]: **0.61**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **4.3**

Temperature indicator value: **5.6**

Moisture indicator value: **4**

Reaction indicator value: **8**

Nutrient indicator value: **3.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.16**

Disturbance frequency (herb layer): **1.29**

Disturbance severity: **0.61**

Disturbance severity (herb layer): **0.18**

Mowing frequency: **0.06**

Grazing pressure: **0.21**

Soil disturbance: **0.09**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) \*Carpino-Fagetea sylvaticae\*](#),  
[CB \(PUB\) \*Quercetea pubescentis\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Constant species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**

Continentality amplitude: **2**