

Frangula rupestris

Habitus and growth type

Plant height [m]: **0.58**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **7**

Moisture indicator value: **3**

Reaction indicator value: **7**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.14**

Disturbance frequency (herb layer): **1.29**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.17**

Mowing frequency: **0.05**

Grazing pressure: **0.24**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#), [CD \(RHA\) Crataego-Prunetea](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**