Fumana laevipes

Habitus and growth type

Plant height [m]: **0.29**
Life span: **Perennial**
Life form: **Chamaephyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **8**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.15**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **9.3**
Moisture indicator value: **3**
Reaction indicator value: **8**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.45**
Disturbance frequency (herb layer): **1.39**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.08**
Grazing pressure: **0.33**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz