Fumana procumbens

Habitus and growth type

Plant height [m]: **0.15**
Life span: **Perennial**
Life form: **Chamaephyte**
Spinescence: **not spinescent**

Leaf

Leaf type (woody plants only): **Needle-like**
Leaf phenology (woody plants only): **Evergreen**
Specific leaf area [mm^2/mg]: **8.22**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **2.8**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.9**
Temperature indicator value: **7**
Moisture indicator value: **1.9**
Reaction indicator value: **8.4**
Nutrient indicator value: **1.4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.12**
Disturbance frequency (herb layer): **1.73**
Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.4**
Mowing frequency: **0.34**
Grazing pressure: **0.38**
Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***, **FP (ONO) *Festuco hystricis-Ononidetea striatae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe**, **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**, **S73 Western Mediterranean mountain hedgehog-heath**
Constant species of EUNIS habitats: **R16 Perennial rocky grassland of Central and South-Eastern Europe**, **R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe**, **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**, **R1H Iberian oromediterranean basiphilous dry grassland**, **S61 Western basiphilous garrigue**, **S73 Western Mediterranean mountain hedgehog-heath**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz