

# ***Fumaria bastardii***

## **Habitus and growth type**

Plant height [m]: **0.59**

Life span: **Annual**

Life form: **Therophyte**

## **Flower**

Flowering period: **February-May**

## **Fruit, seed and dispersal**

Seed mass [mg]: **2.53**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **8.5**

Moisture indicator value: **3.3**

Reaction indicator value: **6**

Nutrient indicator value: **6.5**

Salinity indicator value: **0**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [\*\*QC \(CHE\) Chenopodietea\*\*](#)

## Broad habitat

Occurrence in broad habitats: **Scrub, Synanthropic**