

# ***Fumaria densiflora***

## **Habitus and growth type**

Plant height [m]: **0.61**

Life span: **Annual**

Life form: **Therophyte**

## **Flower**

Flowering period: **February-April**

## **Fruit, seed and dispersal**

Seed mass [mg]: **2.8**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **8.3**

Temperature indicator value: **8.3**

Moisture indicator value: **3**

Reaction indicator value: **6.7**

Nutrient indicator value: **6**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **1.16**

Disturbance frequency (herb layer): **2.06**

Disturbance severity: **0.87**

Disturbance severity (herb layer): **0.83**

Mowing frequency: **0.55**

Grazing pressure: **0.17**

Soil disturbance: **0.66**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**