

Fumaria schleicheri

Habitus and growth type

Plant height [m]: **0.22**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **April-October**

Fruit, seed and dispersal

Seed mass [mg]: **1.98**

Dispersal mode: **Myrmecochory, Anthropochory**

Dispersal distance class: **3, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **6.9**

Moisture indicator value: **4**

Reaction indicator value: **7.4**

Nutrient indicator value: **6.4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.86**

Disturbance frequency (herb layer): **1.69**

Disturbance severity: **0.75**

Disturbance severity (herb layer): **0.65**

Mowing frequency: **0.33**

Grazing pressure: **0.18**

Soil disturbance: **0.55**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

Broad habitat

Occurrence in broad habitats: **Forest, Synanthropic**

Distribution

Continentality: **6**

Continentality amplitude: **5**