

# *Gagea soleirolii*

## Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **4**

Moisture indicator value: **4**

Reaction indicator value: **3**

Nutrient indicator value: **3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.43**

Disturbance frequency (herb layer): **1.63**

Disturbance severity: **0.33**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.03**

Grazing pressure: **0.28**

Soil disturbance: **0.13**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FM \(PIL\) \*Saginetea piliferae\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#), [R1N Open Iberian supramediterranean dry acid and neutral grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**