

# *Galatella villosa*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **1.37**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8.8**

Temperature indicator value: **6.3**

Moisture indicator value: **3.4**

Reaction indicator value: **8**

Nutrient indicator value: **2**

Salinity indicator value: **4**

### Disturbance indicator values

Disturbance frequency: **1.95**

Disturbance frequency (herb layer): **2.05**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.48**

Mowing frequency: **0.62**

Grazing pressure: **0.43**

Soil disturbance: **0.2**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#), [R1B Continental dry grassland \(true steppe\)](#), [R1C Desert steppe](#)

Constant species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#), [R1B Continental dry grassland \(true steppe\)](#), [R1C Desert steppe](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**