

Galinsoga parviflora

Habitus and growth type

Plant height [m]: **0.34**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **39.08**

Flower

Flowering period: **July-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.24**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**

Temperature indicator value: **6**

Moisture indicator value: **5.5**

Reaction indicator value: **5.5**

Nutrient indicator value: **7.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.74**

Disturbance frequency (herb layer): **1.95**

Disturbance severity: **0.83**

Disturbance severity (herb layer): **0.83**

Mowing frequency: **0.56**

Grazing pressure: **0.14**

Soil disturbance: **0.76**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) *Papaveretea rhoeadis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V11 Intensive unmixed crops](#), [V12 Mixed crops of market gardens and horticulture](#)

Constant species of EUNIS habitats: [V11 Intensive unmixed crops](#), [V12 Mixed crops of market gardens and horticulture](#), [V15 Bare tilled, fallow or recently abandoned arable land](#)

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**,

Synanthropic

Distribution

Continentality amplitude: **5**