

Galium aparine

Habitus and growth type

Plant height [m]: **0.92**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **30.76**

Flower

Flowering period: **January-September, December**

Fruit, seed and dispersal

Seed mass [mg]: **9.42**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.6**

Temperature indicator value: **5.7**

Reaction indicator value: **6.3**

Nutrient indicator value: **7.6**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.44**

Disturbance frequency (herb layer): **1.43**

Disturbance severity: **0.67**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.23**

Grazing pressure: **0.17**

Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [HA \(POP\) *Alno glutinosae-Populetea albae*](#), [QG \(EPI\) *Epilobietea angustifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N1E Black Sea broad-leaved coastal dune forest](#)

Constant species of EUNIS habitats: [N1A Atlantic and Baltic coastal dune scrub](#), [N1D Atlantic and Baltic broad-leaved coastal dune forest](#), [N1E Black Sea broad-leaved coastal dune forest](#), [R55 Lowland moist or wet tall-herb and fern fringe](#), [R57 Herbaceous forest clearing vegetation](#), [S32 Temperate Rubus scrub](#), [S35 Temperate and submediterranean thorn scrub](#), [S37 *Corylus avellana* scrub](#), [S38 Temperate forest clearing scrub](#), [S91 Temperate riparian scrub](#), [S93 Mediterranean riparian scrub](#), [T11 Temperate Salix and Populus riparian forest](#), [T12 *Alnus glutinosa*-*Alnus incana* forest on riparian and mineral soils](#), [T13 Temperate hardwood riparian forest](#), [T14 Mediterranean and Macaronesian riparian forest](#), [T1E *Carpinus* and *Quercus* mesic deciduous forest](#), [T1G *Alnus cordata* forest](#), [T1H Broadleaved deciduous plantation of non site-native trees](#), [T25 Phoenix theophrasti vegetation](#), [V11 Intensive unmixed crops](#), [V12 Mixed crops of market gardens and horticulture](#), [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#), [V15 Bare tilled, fallow or recently abandoned arable land](#), [V33 Dry Mediterranean land with unpalatable non-vernal herbaceous vegetation](#), [V39 Mesic perennial anthropogenic herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **5**