

# *Galium fruticosum*

## Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.25**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.28**

Disturbance frequency (herb layer): **1.37**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.33**

Mowing frequency: **0.1**

Grazing pressure: **0.31**

Soil disturbance: **0.16**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*KC \(ASP\) \*Asplenietea trichomanis\*\*\*](#)

## Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**