

Genista cinerea

Habitus and growth type

Plant height [m]: **0.73**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **8**

Moisture indicator value: **3**

Reaction indicator value: **3**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.29**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.14**

Grazing pressure: **0.29**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) *Quercetea ilicis*](#), [FB \(ROS\) *Ononido-Rosmarinetea*](#), [FE \(CYT\) *Cytisetea scopario-striati*](#), [FP \(ONO\) *Festuco hystricis-Ononidetea striatae*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**