

Genista salzmannii

Habitus and growth type

Plant height [m]: **0.48**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **2.1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **6.5**

Moisture indicator value: **2**

Reaction indicator value: **3**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.69**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.41**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.02**

Grazing pressure: **0.26**

Soil disturbance: **0.09**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FQ \(GEN\) Carici-Genistetea lobelii](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S74 Central Mediterranean mountain hedgehog-heath](#)

Constant species of EUNIS habitats: [S74 Central Mediterranean mountain hedgehog-heath](#)

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**