

Gentiana lutea

Habitus and growth type

Plant height [m]: **0.82**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **14.59**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **1.13**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**

Temperature indicator value: **3.3**

Moisture indicator value: **5.1**

Reaction indicator value: **3.8**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.66**

Disturbance frequency (herb layer): **1.46**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.26**

Grazing pressure: **0.26**

Soil disturbance: **0.14**

Habitat and sociology

EUNIS habitat

Constant species of EUNIS habitats: [R23 Mountain hay meadow](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **3**

Continentality amplitude: **3**