

# *Gentianella campestris*

## Habitus and growth type

Plant height [m]: **0.16**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **29.35**

## Flower

Flowering period: **May-October**

## Fruit, seed and dispersal

Seed mass [mg]: **0.18**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **4.4**

Moisture indicator value: **5.1**

Nutrient indicator value: **2.1**

Salinity indicator value: **0.2**

### Disturbance indicator values

Disturbance frequency: **1.46**

Disturbance frequency (herb layer): **1.78**

Disturbance severity: **0.39**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.23**

Grazing pressure: **0.32**

Soil disturbance: **0.16**

## **Habitat and sociology**

Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

## **Distribution**

Continentality: **3**

Continentality amplitude: **3**