

# *Glandora prostrata*

## Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Spinescence: **not spinescent**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **6.5**

Moisture indicator value: **4**

Reaction indicator value: **2.5**

Nutrient indicator value: **1.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.43**

Disturbance frequency (herb layer): **1.35**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.07**

Grazing pressure: **0.28**

Soil disturbance: **0.12**

## Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S62 Western acidophilous garrigue](#), [T29 Broadleaved evergreen plantation of non site-native trees](#)

Broad habitat

Occurrence in broad habitats: **Heathland, Scrub, Forest**