

Hedysarum boveanum

Habitus and growth type

Plant height [m]: **0.32**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Spinescence: **not spinescent**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **5.9**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **7**

Moisture indicator value: **2.5**

Reaction indicator value: **7.5**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.77**

Disturbance frequency (herb layer): **1.89**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.19**

Mowing frequency: **0**

Grazing pressure: **0.44**

Soil disturbance: **0.19**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**