

Helianthemum salicifolium

Habitus and growth type

Plant height [m]: **0.15**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **7.61**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.28**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **7.8**

Moisture indicator value: **2.7**

Reaction indicator value: **6.3**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.23**

Disturbance frequency (herb layer): **2.06**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.63**

Mowing frequency: **0.3**

Grazing pressure: **0.37**

Soil disturbance: **0.32**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) *Helianthemetea guttati*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#)

Constant species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**