

Helosciadium nodiflorum

Habitus and growth type

Plant height [m]: **0.36**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **61.88**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.42**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7.1**

Moisture indicator value: **10**

Reaction indicator value: **6.8**

Nutrient indicator value: **6.1**

Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **1.14**

Disturbance frequency (herb layer): **1.74**

Disturbance severity: **0.37**

Disturbance severity (herb layer): **0.33**

Mowing frequency: **0.18**

Grazing pressure: **0.2**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OD \(PHR\) *Phragmito-Magnocaricetea*](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **5**