Hieracium sabaudum

Habitus and growth type

Plant height [m]: **0.92**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **45.57**

Flower

Flowering period: **July-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.53**
Dispersal mode: **Anemochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.3**
Temperature indicator value: **6**
Moisture indicator value: **4.8**
Reaction indicator value: **3.8**
Nutrient indicator value: **2.4**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.08**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.34**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.16**
Mowing frequency: **0.06**
Grazing pressure: **0.2**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae*, CC (QUE) *Quercetea robori-petraeae*, CL (GER) *Trifolio-Geranietea sanguinei***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R52 Forest fringe of acidic nutrient-poor soils, T1B Acidophilous Quercus forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Synanthropic**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz