

# *Hierochloa australis*

## Habitus and growth type

Plant height [m]: **0.38**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.95**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **5.9**

Temperature indicator value: **6.2**

Moisture indicator value: **4.1**

Nutrient indicator value: **3.2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.03**

Disturbance frequency (herb layer): **1.13**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0.01**

Grazing pressure: **0.2**

Soil disturbance: **0.11**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) \*Carpino-Fagetea sylvaticae\*](#),  
[CB \(PUB\) \*Quercetea pubescentis\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

## Distribution

Continentality: **6**

Continentality amplitude: **1**