

Honckenya peploides

Habitus and growth type

Plant height [m]: **0.13**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **19.04**

Fruit, seed and dispersal

Seed mass [mg]: **5.52**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8.8**

Temperature indicator value: **5.3**

Moisture indicator value: **5.8**

Reaction indicator value: **7**

Nutrient indicator value: **6.8**

Salinity indicator value: **4.5**

Disturbance indicator values

Disturbance frequency: **2.01**

Disturbance frequency (herb layer): **2.11**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.09**

Grazing pressure: **0.17**

Soil disturbance: **0.71**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [JC \(CAK\) *Cakiletea maritimae*](#), [JE \(HON\) *Honckenyo peploidis-Leymetea arenarii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N11 Atlantic, Baltic and Arctic sand beach](#), [N13 Atlantic and Baltic shifting coastal dune](#), [N21 Atlantic, Baltic and Arctic coastal shingle beach](#)

Constant species of EUNIS habitats: [N11 Atlantic, Baltic and Arctic sand beach](#), [N13 Atlantic and Baltic shifting coastal dune](#), [N21 Atlantic, Baltic and Arctic coastal shingle beach](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle**

Distribution

Continentality amplitude: 7