

# *Hyacinthella pallasiana*

## Habitus and growth type

Life span: **Perennial**

Life form: **Geophyte**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **5.8**

Moisture indicator value: **4**

### Disturbance indicator values

Disturbance frequency: **1.8**

Disturbance frequency (herb layer): **1.8**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.22**

Grazing pressure: **0.37**

Soil disturbance: **0.19**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**