

Hypericum caprifolium

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **7.6**

Moisture indicator value: **7.2**

Reaction indicator value: **6.9**

Nutrient indicator value: **3.9**

Salinity indicator value: **1.1**

Disturbance indicator values

Disturbance frequency: **1.48**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **1.17**

Grazing pressure: **0.31**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R31 Mediterranean tall humid inland grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**