

Hypochaeris cretensis

Habitus and growth type

Plant height [m]: **0.25**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.43**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Moisture indicator value: **5**

Nutrient indicator value: **3.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.88**

Disturbance frequency (herb layer): **1.71**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.16**

Grazing pressure: **0.38**

Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FH \(BUL\) *Poetea bulbosae*](#), [FQ \(GEN\) *Carici-Genistetetea lobelii*](#), [KC \(ASP\) *Asplenietea trichomanis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#), [S74 Central Mediterranean mountain hedgehog-heath](#)

Constant species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**