

Chaerophyllum villarsii

Habitus and growth type

Plant height [m]: **0.75**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **6.1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **2.9**

Moisture indicator value: **6.3**

Reaction indicator value: **6.5**

Nutrient indicator value: **7.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.22**

Disturbance frequency (herb layer): **1.36**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.07**

Grazing pressure: **0.19**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CS \(MUL\) *Mulgedio-Aconitetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R23 Mountain hay meadow](#), [S25 Subalpine and subarctic deciduous scrub](#)

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**

Continentality amplitude: **2**