Chenopodium opulifolium

Habitus and growth type

Plant height [m]: **0.66**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **July-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.59**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.9**
Moisture indicator value: **4.4**
Reaction indicator value: **7**
Nutrient indicator value: **6.5**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.81**
Disturbance frequency (herb layer): **2.01**
Disturbance severity: **0.86**
Disturbance severity (herb layer): **0.85**
Mowing frequency: **0.95**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.11**
Soil disturbance: **0.75**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QB (SIS) *Sisymbrietea***

Broad habitat

Occurrence in broad habitats: **Synanthropic**

Distribution

Continentality: **6**
Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz