Chenopodium suecicum

Habitus and growth type

Plant height [m]: **0.62**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.82**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Temperature indicator value: **5.7**
Moisture indicator value: **4.6**
Reaction indicator value: **6.2**
Nutrient indicator value: **7.2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.04**
Disturbance frequency (herb layer): **2.18**
Disturbance severity: **0.86**
Disturbance severity (herb layer): **0.86**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **1.04**
Grazing pressure: **0.12**
Soil disturbance: **0.78**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QB (SIS) *Sisymbrietea***

Broad habitat

Occurrence in broad habitats: **Synanthropic**

Distribution

Continentality: **6**
Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz