

Chenopodium suecicum

Habitus and growth type

Plant height [m]: **0.62**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.82**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **5.7**

Moisture indicator value: **4.6**

Reaction indicator value: **6.2**

Nutrient indicator value: **7.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.04**

Disturbance frequency (herb layer): **2.18**

Disturbance severity: **0.86**

Disturbance severity (herb layer): **0.86**

Mowing frequency: **1.04**

Grazing pressure: **0.12**

Soil disturbance: **0.78**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QB \(SIS\) *Sisymbrietea*](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**

Distribution

Continentality: **6**

Continentality amplitude: **6**