Chrysosplenium oppositifolium

Habitus and growth type

Plant height [m]: **0.09**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **23.35**

Fruit, seed and dispersal

Seed mass [mg]: **0.07**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.2**
Temperature indicator value: **5**
Moisture indicator value: **9.3**
Reaction indicator value: **5.4**
Nutrient indicator value: **5.2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.36**
Disturbance frequency (herb layer): **1.43**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.27**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.04**
Grazing pressure: **0.14**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OA (MON) *Montio-Cardaminetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **P2N Spring**
Constant species of EUNIS habitats: **P2N Spring**
Dominant species of EUNIS habitats: **P2N Spring**

Broad habitat

Occurrence in broad habitats: **Spring, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **2**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz